




LIFE TECHNICAL GUIDE - 01

TG01-0.0-Europe

Version DRAFT 0.0 - Europe - English

(January/2021)

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 14/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 2 of 44 |

OBJECTIVE

To establish the minimum performance that each organization/producer must achieve in biodiversity and ecosystem services conservation actions, considering its size and impact.

APPLICATION

This document applies to organizations and producers who want to contribute to the conservation of biodiversity and ecosystem services and require a support tool to assess and monitor their impacts and their minimum performance for conservation.


For LIFE certified organizations in previous versions, this document will become effective from their first follow-up audit after its publication. For other organizations/producers, this document will automatically apply from the publication date.

APPROVAL

Document approved by the LIFE Institute Board of Directors.

Rights reserved by copyright law in Brazil and abroad according to the terms set forth in Brazilian and foreign legislations relevant to the subject. Any kind of reproduction of this document or part of its content requires the express written permission by the LIFE Institute.

LIFE Institute
 Rua Victor Benato, 210 Bosque Zaninelli, UNILIVRE, Pilarzinho
 CEP: 82120-110 – Curitiba – PR
 Tel: +55 41 3253-7884
www.institutolife.org

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 3 of 44 |

LIST OF ACRONYMS

LC_O: Land cover of the organization/producer

LC_E: Ecoregion original land cover

BMP: Biodiversity Minimum Performance

BPI: Biodiversity Pressure Index

BPP: Biodiversity Positive Performance

boe: Barrel of oil equivalent

DAB_{CHR}: Demand-Availability Balance of the EU most critical country

DAB_{OHR}: Demand-Availability Balance of the country where the organization/producer is located

EC: Total amount of energy consumed

EC_i: Energy consumption from source *i*

EG: Total amount of greenhouse gas emissions

EG_i: Amount of greenhouse gas emitted *i*

EU: Union European

GHG: Greenhouse Gases

GWP: Global warming potential

ID_i: Impact of type *i* waste destination

ID_{max}: Maximum impact observed among the types of waste destination

IE_i: Impact of energy source *i*

IE_{max}: Maximum impact observed among energy sources

PI_i: Pressure index of aspect *i*


PV: Pressure value

IPCC: Intergovernmental Panel on Climate Change

LIFE-EU-TG02: LIFE Technical Guide 02

m³/s: Cubic meter per second

m³: Cubic meter

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 4 of 44 |

NIS: National Interconnected System

APBE: Action Plan for Biodiversity and Ecosystem Services

QSV_{GHG}: Quantity and severity value for the greenhouse gases aspect

QV: Quantity value

RV: Reference value

SV: Severity value

tCO_{2e}: Ton of CO₂ equivalent

Tep (toe): Ton of Oil Equivalent

EU: European Union

WG: Total amount of hazardous and non-hazardous waste generated

WG_i: Waste generated of type *i*

WU: Total amount of water used

WWF: World Wide Fund for Nature

DRAFT



| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 14/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 5 of 44 |

TABLE OF CONTENTS

| | |
|--|-----------|
| 1. INTRODUCTION | 6 |
| 2. BIODIVERSITY PRESSURE INDEX (BPI) | 7 |
| 2.1 CALCULATION OF THE BIODIVERSITY PRESSURE INDEX (BPI)..... | 8 |
| 2.1.1 Quantity and Severity Values | 8 |
| 2.1.2 Pressure Values | 10 |
| 2.1.3 Pressure Indexes..... | 10 |
| 2.1.4 Calculation of the Biodiversity Pressure Index | 11 |
| 2.2 DATA REQUIRED TO CALCULATE THE BPI..... | 11 |
| 2.2.1 Waste Generation..... | 12 |
| 2.2.2 Water Consumption..... | 14 |
| 2.2.3 Energy Consumption..... | 15 |
| 2.2.4 Land cover | 15 |
| 2.2.5 Emission of Greenhouse Gases..... | 16 |
| 3. BIODIVERSITY MINIMUM PERFORMANCE (BMP) | 17 |
| 4. FLOW CHARTS FOR CALCULATING BPI AND BMP | 18 |
| 5. REFERENCES | 19 |
| 6. GLOSSARY | 21 |
| 7. APPENDIX..... | 22 |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 14/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 6 of 44 |

1. INTRODUCTION

The Biodiversity Pressure Index (BPI) is an index developed by the LIFE Institute to define, compare and monitor, in a same scale, the impact of any organization/producer to biodiversity and ecosystem services, serving as an important management tool.


From the calculation of the BPI, organizations and producers who wish to contribute to biodiversity may know and carry out the minimum performance in conservation actions that would be more appropriate to their size and impact.

This document introduces the concept and the manner of obtaining the Biodiversity Pressure Index (BPI) and the minimum performance in biodiversity (BMP) relating to every size and impact.

In addition to the use of these tools for public and private management, organizations and producers who achieve or exceed the minimum performance set, can request a third-party assessment so as to obtain an external recognition on their performance in favor of biodiversity. In this case, LIFE methodology can be granted whenever an organization/producer:

- ✓ Achieves a performance in biodiversity conservation equal to or higher than the minimum set according to the methodology herein described. This positive performance must be demonstrated through an Action Plan for Biodiversity and Ecosystem Services (APBE), assessed and rated according to the document LIFE-EU-TG02.
- ✓ Meets the minimum indicators for biodiversity management described in LIFE Methodology Standards (LIFE-EU-CS).

This document applies to industry, services, and the primary sector (farming areas: agriculture, forestry, animal production and aquaculture), whereas it does not apply to extractivism activities.

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 7 of 44 |

2. BIODIVERSITY PRESSURE INDEX (BPI)

Aiming to establish a metric for scaling and comparing impacts to biodiversity, making it possible to define relative performance for conservation, the Biodiversity Pressure Index (BPI) was developed.

The 5 environmental aspects measured and assessed to calculate the BPI were selected from the performance of public meetings for the definition and selection of relevant variables for the index, both for their relation to the main causes of global biodiversity loss¹ and for their data collection in organizations of any size and sector.

As a result of this analysis, we selected those aspects that had higher viability and ease of data collection and direct relationship with official data available: waste generation; water usage; energy consumption; land cover; greenhouse gas emission.

The BPI is obtained through information relative to the quantity and severity relating to these 5 selected environmental aspects.

Information on the quantity of environmental aspects assessed, or “Quantity Value”, refers to a direct relationship between the data of the organization/producer compared to an official data for this aspect in the European Union. This comparison generates a quantity value of impact for each environmental aspect referring to its contribution to the regional total.


Information on severity, or “Severity Value”, considers specific information for each environmental aspect, which allows to define their criticality: water availability in the region, potential for global warming from the gases emitted, impact of the energy sources used, health hazard, the disposal of waste generated by the activities, and national fragility of the ecoregion occupied by the enterprise. This information, although qualitative, is quantitatively represented by the severity values, which range between 0 and 1 and may be called severity factors.

By multiplying the quantity values of impact by their severity factors, “Pressure Values” (PV) are generated for each environmental aspect. For comparison purposes, these pressure values are transformed into “Pressure Indexes” (PI), with the purpose of being mathematically distributed on the same scale, from zero to one thousand. This distribution has as reference the value of greatest impact known in the region for each environmental aspect.

The simple average of the Pressure Indexes (PI) for each one of the environmental aspects, results in the Biodiversity Pressure Index (BPI).

The following sections of the document present the steps for calculating the BPI and the required Information from organizations and producers for its calculation.


¹ Destruction of habitats; climate changes; introduction of invasive exotic species; over-exploitation of species; pollution (*Millennium Ecosystem Assessment*, 2005).

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 8 of 44 |

2.1 CALCULATION OF THE BIODIVERSITY PRESSURE INDEX (BPI)

This section of the document introduces the steps and equations used to calculate the BPI.

FIGURE 1. Steps for calculating the Biodiversity Pressure Index (BPI).




2.1.1 Quantity and Severity Values

Table 1 presents the equations used to calculate the quantity and severity values for each environmental aspect.

Table 1 - Equations of Quantity Values (QV) and Severity Values (SV) to calculate the BPI for each environmental aspect

| ENVIRONMENTAL ASPECT | QUANTITY | SEVERITY |
|--------------------------|--|---|
| Waste Generation | $QV_{\text{WASTE}} = \frac{WG}{RV_{\text{WASTE}}}$ | $SV_{\text{WASTE}} = \frac{\sum_{i=1}^n (\%WG_i \times ID_i)}{ID_{\text{max}}}$ |
| Water Consumption | $QV_{\text{WATER}} = \frac{WU}{RV_{\text{WATER}}}$ | $SV_{\text{WATER}} = \frac{DAB_{\text{OHR}}}{DAB_{\text{CHR}}}$ |


| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 9 of 44 |

| ENVIRONMENTAL ASPECT | QUANTITY | SEVERITY |
|----------------------------------|--|--|
| Energy Consumption | $QV_{ENERGY} = \frac{EC}{RV_{ENERGY}}$ | $SV_{ENERGY} = \frac{\sum_{i=1}^n (\%EC_i \times IE_i)}{IE_{max}}$ |
| Occupation of Natural Land Cover | $QV_{Land\ cover} = \frac{LC_O}{LC_E}$ | $SV_{Land\ Cover} = \frac{E_{if}}{100}$ |
| Emission of Greenhouse Gases | $QSV_{GHG} = \left(\frac{\sum_{i=1}^n (GE_i \times GWP_i)}{RV_{GHG}} \right)$ | |

Table 2 describes the terms that make up the equations presented in Table 1.

Table 2 – Terms used in the equations for the quantity and severity values

| EQUATION | TERMS USED |
|--------------------|--|
| QV_{WASTE} | <p>QV_{WASTE}= Quantity Value for Waste</p> <p>WG= Total quantity of hazardous and non-hazardous waste generated by the organization/producer (t/year)</p> <p>RV_{WASTE}= Reference Value for waste (t/year) according to Appendix.</p> |
| QV_{WATER} | <p>QV_{WATER}= Quantity Value for Water</p> <p>WU= Consumption of water used by the organization/producer (m³/year)</p> <p>RV_{WATER}= Reference Value for water (m³/year), according to Appendix.</p> |
| QV_{ENERGY} | <p>QV_{ENERGY}= Quantity Value for Energy</p> <p>EC= Total quantity of energy consumed by the organization/producer (toe/year)</p> <p>RV_{ENERGY}= Reference Value for Energy (toe/year), according to Appendix.</p> |
| $QV_{LAND\ COVER}$ | <p>$QV_{Land\ Cover}$= Quantity Value for land cover</p> <p>LC_O= Land cover of the organization/producer (hectares)</p> <p>LC_E= Land cover of the ecoregion in which the organization/producer is located (hectares), according to Appendix</p> |
| QSV_{GHG} | <p>QSV_{GHG}= Quantity and Severity Value for Greenhouse Gases</p> <p>GE_i= Quantity of greenhouse gas emissions <i>i</i> emitted by the organization/producer (tCO₂e/year)</p> <p>GWP_i= Global warming potential of greenhouse gas <i>i</i> according to Appendix.</p> <p>RV_{GHG}= Reference value for greenhouse gases (tCO₂e/year) according to Appendix.</p> |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 10 of 44 |

| EQUATION | TERMS USED |
|-------------------|---|
| SV_{WASTE} | SV_{WASTE} = Severity Value for Waste. WG_i = Percentage of waste generation with type “i” destination. ID_i = Impact of destination “i” (ID) listed in the Appendix. ID_{max} = Maximum impact observed between “i” types of destination. |
| SV_{WATER} | SV_{WATER} = Severity Value for the water aspect. DAB_{CHR} = Demand-Availability Balance DAB_{OHR} = Demand-Availability Balance of the country where the organization/producer is located, listed in the Appendix. |
| SV_{ENERGY} | SV_{ENERGY} = Severity value for the energy aspect. EC_i = Percentage of the energy source type <i>i</i> consumed by the organization/producer. IE_i = Impact of the energy source <i>i</i> consumed by the organization/producer, according to the Appendix. IE_{max} = Maximum impact observed between energy sources according to Appendix. |
| $SV_{LAND COVER}$ | $SV_{Land Cover}$ = Severity value for the land cover aspect. E_{IF} = Ecoregion importance factor, according to Appendix. |

2.1.2 Pressure Values


In Table 3, the equations used to obtain the Pressure Value (PV_i) of each aspect *i* are listed.

Table 3 - Calculation of the pressure value for each environmental aspect

| ENVIRONMENTAL ASPECT | PRESSURE VALUE OF THE ASPECT |
|------------------------------|------------------------------|
| Waste Generation | $PV_i = QV_i \times SV_i$ |
| Water Consumption | |
| Energy Consumption | |
| Land Cover | |
| Emission of Greenhouse Gases | $PV_{GHG} = QSV_{GHG}$ |

2.1.3 Pressure Indexes

The Pressure Values (PV) are transformed into Pressure Indexes (PI), which allow the representation of the pressure of each environmental aspect on the same scale, dimensionless, ranging from zero to 1,000. The Pressure Index (PI) is calculated individually for each environmental aspect by the following equation:

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 11 of 44 |

$$PI_i = \left(1 - \frac{1}{1 + a_i PV_i}\right) \times 1000$$

Where in:

PI = Pressure Index of aspect i

a_i = Correction factor² of aspect i , which allows IP to range between 0 and 1,000

PV = Pressure Value of aspect i

2.1.4 Calculation of the Biodiversity Pressure Index

The Biodiversity Pressure Index is obtained by the simple arithmetic average of the Pressure Indexes (PI) of the five environmental aspects assessed:

$$BPI = \frac{PI_{WATER} + PI_{ENERGY} + PI_{GHG} + PI_{WASTE} + PI_{LAND COVER}}{5}$$


The information herein presented is only a description of the calculations used. Achieving the BPI is facilitated through the use of an automated calculation tool provided by LIFE Institute, upon request.

Information on the reference values used in EU as well as for unit conversion can be found in the Appendix (**Reference Information to calculate the BPI in EU**).

2.2 DATA REQUIRED TO CALCULATE THE BPI

This document section presents the data from the organization/producer that need to be informed to calculate the BPI.

² See details in the Appendix.

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 12 of 44 |

Prior to the calculation, it is necessary to define clearly and objectively which unit is being assessed. This information will be used as reference for the entire assessment process, considering the scope rule for LIFE Methodology.


For situations not provided for in this document or in the support tools mentioned therein, the organization/producer must present its own estimate to the auditor, justifying the data presented during the audit. In case of impossibility of assessing one or more cultures or activities in particular, the auditor may temporarily relieve the organization/producer from inserting the environmental aspect in question into the calculation of the BPI, recording on the audit report the need for effort and follow-up of tools and methodologies over time to collect these data.

2.2.1 Waste Generation

- a) Inform the total amount of waste generated by the organization/producer in tons/year, adding all the following situations:
 - i) Any waste, whether treated or not, forwarded to third parties, whether through donation or sale, for treatment, storage or final elimination
 - ii) Waste send to landfills, own or third party
 - iii) Waste stored, internally or by third parties
 - iv) Household and production waste generated within the property
 - v) Other wastes not receiving internal treatment in the organization/property

The data reported must refer to the total waste generated in all processes - direct and indirect, productive, administrative and from maintenance - as long as performed on the physical site which is being assessed.

- b) There is no need to inform wastes destined internally for:
 - i) Production of biogas;
 - ii) Incineration;
 - iii) Co-processing;
 - iv) Reuse;

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 13 of 44 |

v) Recycling.

All consumption of water, energy and land cover relating to these processes must be informed on the other environmental aspects to calculate the impact of the organization/property.

The auditor may request and assess information on wastes eventually not included in the calculation for the purposes of checking compliance through the evaluation of LIFE Methodology Standards.

c) Inform health hazards of wastes generated in:

- i) Hazardous Waste
- ii) Non-hazardous waste


d) Inform the destination of the waste informed in item (a) in:

- i) Reuse
- ii) Recycling
- iii) Composting
- iv) Landfarming
- v) Co-processing
- vi) Biogas
- vii) Storage
- viii) Incineration
- ix) Landfill with biogas utilization
- x) Landfill

When the destination is different from these categories, the organization/producer may select that with the characteristics closest to the informed destination. In such cases, the auditor must mention and justify this choice in the audit report.

Wastes from agricultural production, even if destined to industry, must be recorded as primary production waste and classified according to the type of destination (e.g.: recycling, co-processing, etc.). If the industry receiving this waste is undergoing assesses by LIFE methodology, this material, in this unit assessed, must be considered as an input and not as waste.

Industrial waste used in agriculture must be informed as “landfarming”, to calculate the impact of the waste from the plant assessed.

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 14 of 44 |

If the value presented is an estimate, due to the absence of previous records, the auditor must assess the coherence of the figures provided and record on the audit report the need to begin periodic controls.


2.2.2 Water Consumption

- a) Inform the volume of consumptive water use³ of all processes, direct and indirect, carried out in the physical unit assessed.
- i) **Primary sector:** inform the sum of the values for “green footprint” (water from precipitation stored in plants, evaporated or transpired) and for “blue footprint” (surface or underground water incorporated into the process).
- Agricultural crops: water consumption estimates for each crop can be obtained through online tool from the Water Footprint Network initiative⁴.
 - Animal production: water consumption estimates can be obtained by extrapolation of the individual consumption per animal/head, including watering, washing, etc.
 - Forestry: water consumption estimates for the *Pinus* and *Eucalyptus* genera can be obtained through the LIFE Key calculation tool⁵. In these cases, it is necessary to inform the area planted with each gender and the location of plantations.
- ii) **Secondary sector:** the organization must inform only the consumptive use of blue water (water collected less the water discarded, either as effluent or process losses).
- iii) **Tertiary sector:** the organization must inform only the consumption of blue water. The consumption of blue water can be informed through consumption records, being possible to discount the return volume to the basin only when this information is available.
- b) Inform the country where the assessed enterprise is located.

³ Non-consumptive uses do not need to be reported, e.g.: aquaculture, hydroelectricity, water for dilution and/or purification of effluents.

⁴ *The Water Footprint Assessment Tool* Available at: <http://waterfootprint.org/en/resources/interactive-tools/water-footprint-assessment-tool/>

⁵ Estimates obtained through the LIFE/IPEF project of Forestry water consumption.

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 15 of 44 |

2.2.3 Energy Consumption


a) Inform the total amount of energy consumed (own or acquired by the business unit). Inform the distribution of consumption by sources used:

- i) Energy from the electricity produced by country (grid)
- ii) Biofuels (ethanol)
- iii) Biofuels (Oils and Biodiesel)
- iv) Biogas
- v) Biomass (wood)
- vi) Biomass (residual)
- vii) Mineral Coal
- viii) Sea Energy
- ix) Wind
- x) Natural Gas
- xi) Geothermal
- xii) Hydroelectricity
- xiii) Non-renewable residual
- xiv) Nuclear
- xv) Oil and derived
- xvi) Solar

2.2.4 Land cover

a) Inform the *land cover* (hectares), distributed according to occupancy classes in accordance with MSA (Mean Species Abundance⁶).

⁶ Mean Species Abundance (MSA) is an indicator that describes the changes in the environment in relation to the original ecosystem. The MSA is an indicator of naturalness or intactability of biodiversity, defined as mean abundance of original species in the land cover in question in relation to their abundance in undisturbed ecosystems. An land cover with an MSA of 100% (1.0) means having a biodiversity similar to the natural situation. An MSA of 0% (0.0) means a 2020 LIFE Institute - All rights reserved. Only documents available on the LIFE Institute website can be considered as Official Version.

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 16 of 44 |

b) Inform the ecoregion in which the organization/producer is located. The organization can define its ecoregion more accurately by entering the location data on the map provided by the LIFE Key tool.

c) In the case of agricultural properties bound to leasing contracts or others, inform only the land covers relating to the contract⁷.

d) External land covers to the assessed properties, bound only to conservation actions, must not be accounted for to calculate the BPI.

2.2.5 Emission of Greenhouse Gases

a) The total amount of emissions of all greenhouse gases;

The organization/producer must inform the Total Emissions of each one of the Greenhouse Gases (tCO₂e/year), considering the Scopes 1+2+3 of the GHG Protocol tool.⁸ More detailed information on the scopes of the GHG Protocol is listed in the Appendix and in the document LIFE-IN-RD003.

The GHG Protocol also has a calculation tool specific for the primary sector⁹. Other tools for the inventory of emissions will be accepted, as long as also using the IPCC (Intergovernmental Panel on Climate Change) guidelines¹⁰.

The BPI assesses the negative impacts to biodiversity for all environmental aspects considered. Thus, for this step, only greenhouse gas emissions will be accounted for, and not carbon sequestration. Carbon fixation projects, validated by a third party¹¹, may score as indirect action for biodiversity conservation (strategic line “G4” - LIFE-EU-TG02).

completely destroyed ecosystem without remaining original species. The relationship of the MSA classes for land cover are in the Appendix. For more details: <http://www.globio.info/background-msa>


⁷ In these cases, legal environmental compliance is mandatory for the entire land cover of the property, even if the contract is bound to a partial area. This mandatory legal compliance must be provided for in contract.

⁸ Cross-sectoral GHG Protocol Tool. Available at <http://twixar.me/sVP>

⁹ GHG Protocol Agricultural Guidelines. Available at <http://twixar.me/cVP>

¹⁰ Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://twixar.me/xVP>

¹¹ Validation by recognized initiatives relative to the topic or by consulting works based in detailed, justified and recognized methodologies.

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 17 of 44 |

3. BIODIVERSITY MINIMUM PERFORMANCE (BMP)

The minimum performance in biodiversity conservation for the LIFE Methodology is determined by two factors: the Biodiversity Pressure Index (BPI) and the company's turnover.

The BMP is obtained through the following equation:

$$BMP = 50 \times BPI^x \times TO$$

Where in:

BMP: Biodiversity Minimum Performance

BPI: Biodiversity Pressure Index


TO: Turnover

x, y: calibration factors of BMP


Once the BMP is calculated the organization has to evaluate and compare it with its Biodiversity Positive Performance (BPP).


Biodiversity Positive Performance (BPP) is related to the score of the organization's Biodiversity and Ecosystem Services Action Plan (BESAP).

The methodology for scoring BPP can be found on LIFE Technical Guide 02.

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 14/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 18 of 44 |

4. FLOW CHARTS FOR CALCULATING BPI AND BMP



| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 19 of 44 |

5. REFERENCES

GLO BIO. **GLO BIO3: A Framework to Investigate Options for Reducing Global Terrestrial Biodiversity Loss.**

2009. Available at <<https://www.globio.info/globio3-framework-to-investigate-options-for-reducing-global-terrestrial-biodiversity-loss>>. Access on: 05 October 2020.

GREEN HOUSE GAS PROTOCOL. **Calculation tool 2017.** Available at: <<https://ghgprotocol.org/calculation-tools>>. Access on: 16 October 2020.

WATER FOOTPRINT NETWORK. **The Water Footprint Assessment Manual** Available at: <http://waterfootprint.org/media/downloads/TheWaterFootprintAssessmentManual_2.pdf>. Access on: 15 Oct. 2015.

WORLD WIDE FUND FOR NATURE (WWF). **Terrestrial ecoregions of the world (TEOW): A new map of life on Earth.** 2001. Available at: <<https://www.worldwildlife.org/publications/terrestrial-ecoregions-of-the-world>>. Access on: 7 September 2020

EUROSTAT. Generation of waste by economic activity. Available at: <https://ec.europa.eu/eurostat/databrowser/product/view/ENV_WASGEN> Access on: 16 October 2020.

EUROSTAT. Final energy consumption by product. 2017. Available at: <https://ec.europa.eu/eurostat/databrowser/product/view/NRG_BAL_S> Access on: 16 October 2020.

EUROPEAN ENVIRONMENTAL AGENCY. Total GHG Emissions in EU-27 . 2017. Available at:<<https://www.eea.europa.eu/data-and-maps/data/data-viewers/greenhouse-gases-viewer>> Access on: 16 October 2020.


WORLDOMETER. Water use by country 2017. Available at:<<https://www.worldometers.info/water/>> Access on: 16 October 2020.

INDEX MUNDI. Malta - Renewable internal freshwater resources. 2017. Available at:<<https://www.indexmundi.com/facts/malta/indicator/ER.H2O.INTR.K3>> Access on: 16 October 2020.

BIRDLIFE INTERNATIONAL. World Database of Key Biodiversity Areas. 2020 Available at: <<http://www.keybiodiversityareas.org/site/requestgis>>. Access on: 7 September 2020.

EUROPEAN ENVIRONMENT AGENCY. Natura 2000 End 2019 – Shapefile. Available at: <<https://www.eea.europa.eu/data-and-maps/data/natura-11/natura-2000-spatial-data/natura-2000-shapefile-1>> Access on: 7 September 2020.


EUROPEAN ENVIRONMENT AGENCY. Distribution of habitats, 2019, vector polygon. Available at: <<https://www.eea.europa.eu/data-and-maps/data/article-17-database-habitats-directive-92-43->

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 20 of 44 |

eec/distribution-of-species-zipped-shapefile-vector-polygon/distribution-of-habitats-zipped-shapefile-vector-polygon> Access on: 7 September 2020.

CONSERVATION INTERNATIONAL. Biodiversity Hotspots Revisited. 2011. Available at: <<https://www.arcgis.com/home/item.html?id=e5a7d024c4674cc185f78a99071feb07>> Access on: 7 September 2020.


DRAFT

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 21 of 44 |

6. GLOSSARY

The terms used in this document are available in the LIFE Glossary.

DRAFT

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 22 of 44 |

7. APPENDIX

1. Factor a_i


Factor a_i is the Correction Factor of distribution scale of the Pressure Indexes. The correction factors are determined nationally, aimed at establishing a distribution scale of the impacts from the higher values for each individual impact (productive unit) in the country. In each country, the factor is set so that the maximum value observed for the environmental aspect is equivalent to the value of 950 in a scale from 0 to 1,000.

The Correction Factors presently used in Union European are: **(i) Waste: 1.804.835; (ii) Water: 1.925.155; (iii) Energy: 76.931; (iv) Land Cover: 5.326; (v) Greenhouse Gases: 1.971.**

2. Calibration factors of BMP

The factors of equation BMP are the ones that adjust the regions's conservation performance according to the current practices of organizations, so that all enterprises seek to achieve the best practices. Current practices of organizations in conservation are researched and assessed by local experts.


Calibration factors of BMP in Europe: x) 0,42; y) 0,29.

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 23 of 44 |

3. Reference Values (RV) for environmental aspects

The Reference Value (RV) represents the whole, in terms of quantity, national impact in one year.

| ASPECT | REFERENCE VALUE (RV) | DOCUMENT | YEAR | BASE YEAR | INFORMATION USED |
|---------------|---|--|------|-----------|---|
| WASTE | 2.260.515.093,00 t/year | Eurostat Database | 2020 | 2016 | Estimated generation of total urban solid waste in the European Community. |
| GASES | 3.977.716.368,85 tCO ₂ e/year | Data viewer on greenhouse gas emissions and removals, sent by countries to UNFCCC and the EU Greenhouse Gas Monitoring Mechanism (EU Member States). | 2020 | 2017 | Total greenhouse gas emissions in the UE in CO ₂ e converted through the GWP metric. |
| ENERGY | 640.200.000,00 toe/year | Eurostat Database | 2020 | 2017 | Total energy supply in the European Union (EU) *Except Malta. |
| WATER | 195.047.000.000,00 m ³ /year | Eurostat Database | 2018 | 2017 | Demand for water that corresponds to the flow of withdrawal, that is, water collected intended to various consumption uses. |
| | | WordoMeter Database | 2017 | 2017 | |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 24 of 44 |

4. References for the calculation of the Severity Valor


a) Impact of Destination (ID) of non-hazardous waste generated by the organization

| ASPECT | Reduction of the volume of waste to be disposed in a landfill | Reduction of the potential for contamination of the waste | Generation of new products | Energy reuse | Reduction of the consumption of natural resources | Generation of other waste | Area degradation | Generation of liquid effluents / Possibly contaminated water | Generation of pollutant gases | Sum of the impact | Process score | Severity Index |
|----------------------------------|---|---|----------------------------|--------------|---|---------------------------|------------------|--|-------------------------------|-------------------|---------------|----------------|
| | Positive Impact | | | | | Negative Impact | | | | | | |
| Reuse | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| Recycling | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 2 | 2 | 4 |
| Composting | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 2 | 2 | 4 |
| Landfarming | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 4 | 8 |
| Co-processing | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 3 | 3 | 9 |
| Biogas | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 4 | 3 | 12 |
| Storage | 0 | 0 | 1 | 1 | - | 0 | 0 | 1 | 0 | 3 | 4 | 12 |
| Incineration | 0 | 0 | 1 | 0 | 1 | 1 | - | - | 1 | 4 | 4 | 16 |
| Landfill with biogas utilization | 1 | 1 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 7 | 5 | 35 |
| Landfill | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 9 | 5 | 45 |

Positive Impact: 0 - Presence of positive impact / 1 - Absence of positive impact.

Negative Impact: 0 - Absence of negative impact / 1 - Presence of negative impact

Max ID: 45

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 25 of 44 |

b) Impact of Destination (ID) of hazardous waste generated by the organization

| ASPECT | Reduction of the volume of waste | Reduction of the potential for contamination of the waste | Generation of new products | Energy reuse | Reduction of the consumption of natural resources | Generation of other waste | Area degradation | Generation of liquid effluents / Possibly contaminated water bodies | Generation of pollutant gases | Flammability | Corrosivity | Reactivity | Toxicity | Pathogenicity | Sum of the impact | Process score | Severity Index |
|----------------------------------|----------------------------------|---|----------------------------|--------------|---|---------------------------|------------------|---|-------------------------------|--------------|-------------|------------|----------|---------------|-------------------|---------------|----------------|
| | Positive Impact | | | | | Negative Impact | | | | | | | | | | | |
| Reuse | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 10 | 10 | 10 | 10 | 50 | 0 | 50 |
| Recycling | 0 | 0 | 0 | 0 | 0 | 10 | 0 | 10 | 0 | 10 | 10 | 10 | 10 | 10 | 70 | 1 | 70 |
| Landfarming | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 70 | 1 | 70 |
| Co-processing | 0 | 0 | 0 | 0 | 0 | 10 | 0 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 80 | 1 | 80 |
| Biogas | 0 | 0 | 0 | 10 | 0 | 0 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 90 | 2 | 180 |
| Storage | 0 | 0 | 10 | 10 | - | 0 | 0 | 10 | 0 | 10 | 10 | 10 | 10 | 10 | 80 | 3 | 240 |
| Incineration | 0 | 0 | 10 | 0 | 10 | 10 | - | - | 10 | 10 | 10 | 10 | 10 | 10 | 90 | 4 | 360 |
| Landfill with biogas utilization | 10 | 10 | 0 | 0 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 120 | 5 | 600 |
| Landfill | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 140 | 5 | 700 |


Positive Impact: 0 - Presence of positive impact / 10 - Absence of positive impact.

Negative Impact: 0 - Absence of negative impact / 10 - Presence of negative impact

Max ID: 700

c) Demand-Availability Balance (DAB) by country


| Country | Water availability in the region (m ³ /s) | Water demand in the region (m ³ /s) | Demand-Availability Balance (DAB) |
|----------|--|--|-----------------------------------|
| Austria | 2.473,4 | 110,7 | 0,045 |
| Belgium | 570,8 | 141,4 | 0,248 |
| Bulgaria | 665,9 | 176,4 | 0,265 |
| Croatia | 3.361,2 | 21,7 | 0,006 |
| Cyprus | 31,7 | 7,2 | 0,228 |
| Czechia | 412,2 | 51,8 | 0,126 |
| Denmark | 190,3 | 28,2 | 0,148 |
| Estonia | 412,2 | 54,7 | 0,133 |
| Finland | 3.488,1 | 208,1 | 0,060 |
| France | 6.690,8 | 868,6 | 0,130 |
| Germany | 4.883,3 | 789,0 | 0,162 |
| Greece | 2.156,3 | 324,9 | 0,151 |

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 26 of 44 |

| Country | Water availability in the region (m ³ /s) | Water demand in the region (m ³ /s) | Demand-Availability Balance (DAB) |
|-------------|--|--|-----------------------------------|
| Hungary | 3.297,8 | 136,0 | 0,041 |
| Ireland | 1.648,9 | 24,0 | 0,015 |
| Italy | 6.056,6 | 1084,2 | 0,179 |
| Latvia | 1.109,8 | 6,50 | 0,006 |
| Lithuania | 792,7 | 11,89 | 0,015 |
| Luzembourg | 126,8 | 1,43 | 0,011 |
| Malta | 1,6 | 1,40 | 0,863 |
| Netherlands | 2.885,6 | 288,91 | 0,100 |
| Poland | 1.934,3 | 351,63 | 0,182 |
| Portugal | 2.441,7 | 153,38 | 0,063 |
| Romania | 6.722,5 | 204,65 | 0,030 |
| Slovakia | 1.585,5 | 18,39 | 0,012 |
| Slovenia | 1.014,7 | 29,05 | 0,029 |
| Spain | 3.551,5 | 1015,35 | 0,286 |
| Sweden | 5.517,5 | 75,31 | 0,014 |

d) Severity Value for Water


| Country | $VS_a = BDD(RHO)/BDD(RHC)$ |
|----------|----------------------------|
| Austria | 0,051891608 |
| Belgium | 0,287196970 |
| Bulgaria | 0,307103896 |
| Croatia | 0,007479417 |
| Cyprus | 0,264272727 |
| Czechia | 0,145688811 |
| Denmark | 0,171545455 |
| Estonia | 0,153891608 |
| Finland | 0,069145041 |
| France | 0,150467579 |
| Germany | 0,187283501 |
| Greece | 0,174664773 |
| Hungary | 0,047812500 |
| Ireland | 0,016873689 |
| Italy | 0,207483341 |
| Latvia | 0,006788961 |

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 27 of 44 |

| Country | $VSa = BDD(RHO)/BDD(RHC)$ |
|-------------|---------------------------|
| Lithuania | 0,017386364 |
| Luzembourg | 0,013039773 |
| Malta | 1,000000000 |
| Netherlands | 0,116049201 |
| Poland | 0,210707526 |
| Portugal | 0,072811983 |
| Romania | 0,035286664 |
| Slovakia | 0,013445455 |
| Slovenia | 0,033178977 |
| Spain | 0,331375812 |
| Sweden | 0,015820925 |


e) Participation by energy source in the electricity in EU:

| Participation in the total electricity production by country in the EU (%) | | | | | | |
|--|--------------|---------|-------------------|-------|-------|------------|
| Country | Mineral Coal | Nuclear | Hydro Electricity | Wind | Solar | Geothermal |
| Austria | 24,58 | 0,00 | 56,72 | 10,18 | 0,00 | 8,52 |
| Belgium | 36,33 | 47,56 | 1,26 | 10,55 | 4,06 | 0,23 |
| Bulgaria | 45,77 | 38,99 | 8,32 | 3,26 | 3,53 | 0,13 |
| Croatia | 39,28 | 0,00 | 47,65 | 11,86 | 0,59 | 0,61 |
| Cyprus | 90,71 | 0,00 | 0,00 | 4,88 | 4,41 | 0,00 |
| Cz.Republic | 57,12 | 35,23 | 3,87 | 0,85 | 2,79 | 0,14 |
| Denmark | 39,79 | 0,00 | 0,06 | 56,76 | 3,39 | 0,00 |
| Estonia | 89,03 | 0,00 | 0,28 | 10,69 | 0,00 | 0,00 |
| Finland | 37,36 | 34,71 | 18,60 | 9,06 | 0,27 | 0,00 |
| France | 10,53 | 69,88 | 11,24 | 6,29 | 2,03 | 0,02 |
| Germany | 50,92 | 13,14 | 4,22 | 23,28 | 8,31 | 0,13 |
| Greece | 67,97 | 0,00 | 8,48 | 15,23 | 8,29 | 0,04 |
| Hungary | 44,51 | 48,28 | 0,67 | 2,21 | 4,31 | 0,03 |
| Ireland | 63,65 | 0,00 | 3,88 | 32,46 | 0,00 | 0,00 |
| Italy | 65,81 | 0,00 | 16,54 | 7,07 | 8,57 | 2,00 |
| Latvia | 63,61 | 0,00 | 33,93 | 2,46 | 0,00 | 0,00 |
| Lithuania | 26,77 | 0,00 | 25,14 | 40,27 | 2,02 | 5,80 |
| Luxembourg | 27,61 | 0,00 | 51,17 | 15,08 | 6,14 | 0,00 |
| Malta | 90,25 | 0,00 | 0,00 | 0,00 | 0,00 | 9,75 |
| Netherlands | 82,28 | 3,16 | 0,06 | 9,78 | 4,32 | 0,40 |
| Poland | 87,86 | 0,00 | 1,77 | 9,89 | 0,49 | 0,00 |

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 28 of 44 |

| | | | | | | |
|----------|-------|-------|-------|-------|------|------|
| Portugal | 51,40 | 0,00 | 19,53 | 26,21 | 2,49 | 0,37 |
| Romania | 36,36 | 19,21 | 28,72 | 12,45 | 3,27 | 0,00 |
| Slovakia | 25,06 | 55,32 | 17,16 | 0,01 | 2,26 | 0,19 |
| Slovenia | 30,72 | 37,05 | 30,40 | 0,04 | 1,80 | 0,00 |
| Spain | 42,10 | 21,39 | 10,10 | 20,93 | 5,49 | 0,00 |
| Sweden | 9,49 | 39,12 | 39,29 | 12,10 | 0,00 | 0,00 |

DRAFT


| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 29 of 44 |

f) Impact of energy sources used by the organization (IE)

| IMPACT | | | | | | | | | | | | | |
|-------------------------------------|--------------------------------|-------------------------|---|-----------------------|------------------------|--------------------------------------|--------------------------------------|-----------------------------------|-----------------------------------|------------------------|---|---|------------------------------|
| COMPONENT | WATER | | AIR | | | SOIL | | | | BIOTA | | | IMPACT OF ENERGY SOURCE (IE) |
| ENVIRONMENTAL FACTOR | Water use and / or consumption | Generation of effluents | Emissions of greenhouse gases | Atmospheric emissions | Noise emissions | Movement of soil | | Land use | Generation of solid waste | Occupation of areas | Generation of effluents and solid residues; atmospheric emissions | | |
| POTENTIAL IMPACT | Change in water availability | Change in water quality | Contribution to increased climate warming | Change in air quality | Change in noise levels | Intensification of silting processes | Intensification of erosive processes | Generation of induced earthquakes | Changes in landscape and land use | Change in soil quality | Habitat change and / or reduction | Structural and / or functional change of ecosystems | |
| ENERGY SOURCE | | | | | | | | | | | | | |
| Biofuels (Ethanol) | 9 | 5 | 2 | 5 | 1 | 2 | 5 | n.s | 9 | 1 | 5 | 3 | 47 |
| Biofuel (Oils and Biodiesel) | 9 | 5 | 2 | 5 | 1 | 2 | 5 | n.s | 5 | 5 | 5 | 3 | 47 |
| Biogas | 2 | 1 | 3 | 3 | 1 | n.s | n.s | n.s | 2 | 1 | n.s | n.s | 13 |
| Biomass (wood) | 3 | 1 | 9 | 7 | 3 | 2 | 2 | n.s | 7 | 3 | 9 | 3 | 49 |
| Biomass (residual) | 1 | 1 | 3 | 5 | 1 | 1 | 1 | n.s | 5 | 3 | 1 | 3 | 25 |
| Mineral Coal | 9 | 8 | 10 | 10 | 7 | 9 | 9 | 9 | 10 | 10 | 10 | 9 | 110 |
| Sea Energy | n.s | n.s | n.s | n.s | 2 | n.s | n.s | n.s | 1 | n.s | 5 | 1 | 9 |
| Wind | n.s | n.s | n.s | n.s | 6 | n.s | 1 | n.s | 9 | n.s | 2 | n.s | 18 |
| Natural Gas | 9 | 7 | 9 | 7 | 7 | 4 | 4 | 9 | 9 | 5 | 8 | 6 | 84 |
| Geothermal | 1 | 6 | 1 | 2 | 4 | 1 | 1 | 9 | 9 | 5 | 5 | 1 | 45 |
| Hydroelectricity | 9 | 1 | 1 | 3 | 3 | 10 | 9 | 2 | 10 | 1 | 9 | 1 | 59 |
| Non-renewable residual | 1 | 5 | 10 | 7 | 5 | 1 | 1 | n.s | 5 | n.s | 2 | 1 | 38 |
| Nuclear | 10 | 6 | 1 | 3 | 7 | 9 | 9 | 9 | 10 | 10 | 9 | 5 | 88 |
| Petroleum and byproducts | 9 | 8 | 10 | 10 | 7 | 4 | 4 | 9 | 9 | 8 | 4 | 6 | 88 |
| Solar | 5 | 1 | 1 | n.s | 1 | 1 | 1 | n.s | 6 | 6 | 5 | 5 | 32 |

n.s = not significant

IE_{max}= 110


| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 30 of 44 |

g) Mean Species Abundance (MSA)

| MSA | Classes of Soil Cover |
|------|--|
| 1 | Areas permanently covered with snow or ice considered as undisturbed areas |
| 1 | Areas permanently without vegetation (for example, deserts, high alpine areas) |
| 1 | Minimal disturbance, where flora and fauna species abundance are near pristine |
| 0.7 | Forests with extractive use and associated disturbance like hunting and selective logging, where timber extraction is followed by a long period of re-growth with naturally occurring tree species |
| 0.5 | Areas originally covered with forest or woodlands, where vegetation has been removed, forest is re-growing or has a different cover and is no longer in use |
| 0.2 | Planted forest often with exotic species |
| 1.0 | Grassland or scrubland-dominated vegetation (for example, steppe, tundra, or savannah) |
| 0.7 | Grasslands where wildlife is replaced by grazing livestock |
| 0.1 | Forests and woodlands that have been converted to grasslands for livestock grazing. |
| 0.5 | Agricultural production intercropped with (native) trees. Trees are kept for shade or as wind shelter |
| 0.3 | Subsistence and traditional farming, extensive farming, and low external input agriculture |
| 0.1 | High external input agriculture, conventional agriculture, mostly with a degree of regional specialization, irrigation-based agriculture, drainage-based agriculture. |
| 0.05 | Areas more than 80% built up |

h) Greenhouse Gases and their global warming potential (GWP) for a period of 100 years

| Gas | Chemical formula | GWP |
|--------------------------------------|--|-------|
| Carbon Dioxide | CO ₂ | 1 |
| Methane | CH ₄ | 21 |
| Nitrous Oxide | N ₂ O | 310 |
| Hydrofluorocarbon (HFC) | | |
| HFC-125 | C ₂ HF ₅ | 2,800 |
| HFC-134a | C ₂ H ₂ F ₄ (CH ₂ FCF ₃) | 1,300 |
| HFC-143a | C ₂ H ₃ F ₃ (CF ₃ CH ₃) | 3,800 |
| HFC-152a | C ₂ H ₄ F ₂ (CH ₃ CHF ₂) | 140 |
| Perfluorocarbons (PFC) | | |
| Perfluoromethane (tetrafluoroethane) | CF ₄ | 6,500 |

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 31 of 44 |


| Gas | Chemical formula | GWP |
|-----------------------------------|-------------------------------|--------|
| Perfluorethane (Hexafluoroethane) | C ₂ F ₆ | 9,200 |
| Sulfur hexafluoride | SF ₆ | 23,900 |

Adapted from: Second National Communication of Brazil to the United Nations Framework Convention on Climate Change - Volume 1.

i) Ecoregions of Union European

a) Original land cover and priority conservation of ecoregions in Union European

| Ecoregion | Original land cover (ha) | Overlapping of land cover conservation priority database | Ecoregion Importance Factor (E_{if}) |
|---|--------------------------|--|--|
| Crete Mediterranean forests | 770.378,67 | 770.378,67 | 100,00 |
| Cyprus Mediterranean forests | 525.449,09 | 525.449,09 | 100,00 |
| Iberian conifer forests | 3.446.101,73 | 3.446.101,73 | 100,00 |
| Mediterranean acacia-argania dry woodlands and succulent thickets | 232.534,43 | 232.534,43 | 100,00 |
| Mediterranean woodlands and forests | 2.458,00 | 2.458,00 | 100,00 |
| Southeastern Iberian shrubs and woodlands | 268.168,71 | 268.168,71 | 100,00 |
| Tyrrhenian-Adriatic Sclerophyllous and mixed forests | 7.824.272,49 | 7.824.272,49 | 100,00 |
| Azores temperate mixed forests | 218.074,60 | 218.074,60 | 100,00 |
| Corsican montane broadleaf and mixed forests | 363.364,55 | 363.364,54 | 100,00 |
| South Appenine mixed montane forests | 1.309.479,52 | 1.309.478,39 | 100,00 |
| Canary Islands dry woodlands and forests | 465.859,22 | 465.858,76 | 100,00 |
| Madeira evergreen forests | 74.164,73 | 74.164,52 | 100,00 |
| Southwest Iberian Mediterranean sclerophyllous and mixed forests | 7.008.156,14 | 7.008.126,53 | 100,00 |
| Aegean and Western Turkey sclerophyllous and mixed forests | 7.635.681,86 | 7.635.555,52 | 100,00 |
| Pindus Mountains mixed forests | 2.315.927,04 | 2.315.862,58 | 100,00 |
| Iberian sclerophyllous and semi-deciduous forests | 29.789.114,72 | 29.786.998,49 | 99,99 |
| Northwest Iberian montane forests | 5.740.590,75 | 5.738.649,75 | 99,97 |
| Northeastern Spain and Southern France Mediterranean forests | 8.932.153,37 | 8.925.233,65 | 99,92 |
| Illyrian deciduous forests | 1.664.068,52 | 1.662.407,29 | 99,90 |
| Italian sclerophyllous and semi-deciduous forests | 10.060.224,66 | 10.049.188,80 | 99,89 |
| Euxine-Colchic broadleaf forests | 15.070,95 | 14.592,22 | 96,82 |
| Dinaric Mountains mixed forests | 1.659.958,55 | 1.054.365,37 | 63,52 |
| Rodope montane mixed forests | 2.966.659,23 | 1.766.788,75 | 59,55 |
| Pyrenees conifer and mixed forests | 2.543.322,89 | 1.480.773,47 | 58,22 |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 32 of 44 |

| Ecoregion | Original land cover (ha) | Overlapping of land cover conservation priority database | Ecoregion Importance Factor (E_{if}) |
|--|--------------------------|--|--|
| Scandinavian Montane Birch forest and grasslands | 5.234.378,86 | 2.759.710,51 | 52,72 |
| Appenine deciduous montane forests | 1.614.718,12 | 829.641,15 | 51,38 |
| Carpathian montane forests | 9.139.307,21 | 3.806.743,47 | 41,65 |
| Pontic steppe | 2.435.051,96 | 918.656,42 | 37,73 |
| Alps conifer and mixed forests | 12.539.119,06 | 3.796.115,15 | 30,27 |
| Cantabrian mixed forests | 7.887.524,26 | 2.238.174,00 | 28,38 |
| Baltic mixed forests | 11.022.013,33 | 2.830.999,34 | 25,68 |
| Balkan mixed forests | 12.430.536,13 | 3.093.135,31 | 24,88 |
| Pannonian mixed forests | 25.991.173,98 | 6.394.192,91 | 24,60 |
| Scandinavian and Russian taiga | 57.858.526,35 | 14.166.179,81 | 24,48 |
| North Atlantic moist mixed forests | 1.553.126,30 | 310.084,52 | 19,97 |
| Western European broadleaf forests | 47.489.202,61 | 8.576.189,73 | 18,06 |
| Central European mixed forests | 36.998.830,89 | 6.563.864,09 | 17,74 |
| Lake | 573.300,62 | 72.967,21 | 12,73 |
| Atlantic mixed forests | 38.965.825,62 | 4.534.777,22 | 11,64 |
| Celtic broadleaf forests | 5.136.012,88 | 582.234,84 | 11,34 |
| Po Basin mixed forests | 4.193.635,42 | 443.674,75 | 10,58 |
| Sarmatic mixed forests | 26.897.165,98 | 2.684.084,06 | 9,98 |
| East European forest steppe | 1.975.922,86 | 162.610,93 | 8,23 |


Source: LIFE Institute. Ecoregion Importance Factor calculated through the overlapping of 4 conservation priority area european database: The habitat's Directive (Directive 92/43/EEC), Key Biodiversity Areas (KBA's), Natura 2000 and Biodiversity Hotspots (Conservation International).

5. Factors for Unit Conversation : ADAPTATION BY ECOACSA

a) Relations between Units

| Exponential | Equivalence | Practical relations |
|-----------------------|---|----------------------------|
| (k) kilogram = 10^3 | 1 m ³ = 6.28981 barrels | |
| (M) mega = 10^6 | 1 barrel = 0.158987 m ³ | 1 toe year = 7.2 boe year |
| (G) giga = 10^9 | 1 joule = 0.239 cal | 1 boe year = 0.14 toe year |
| (T) tera = 10^{12} | 1 Btu = 252 cal | 1 toe year = 0.02 boe day |
| (P) peta = 10^{15} | 1 m ³ of oil = 0.872 t (in 1994) | 1 boe day = 50 toe year |
| (E) exa = 10^{18} | 1 toe = 10,000 Mcal | |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 33 of 44 |

b) Coefficients of Caloric Equivalence

| Multiplied by from | to | (m ³) | (1,000 m ³) | (t) | (m ³) | (t) | (t) |
|-----------------------|-------------------------|-------------------|-------------------------|--------------------|-------------------|----------|----------|
| | | Fuel oil | Dry natural gas | Mineral Coal 5,200 | LPG | Firewood | Charcoal |
| Mineral Coal 5,200 | (t) | 0.52 | 0.56 | 1.00 | 0.80 | 1.58 | 0.76 |
| Charcoal | (t) | 0.67 | 0.73 | 1.31 | 1.05 | 2.06 | 1.00 |
| Dry natural gas | (1,000 m ³) | 0.92 | 1.00 | 1.78 | 1.43 | 2.80 | 1.36 |
| LPG | (m ³) | 0.64 | 0.70 | 1.25 | 1.00 | 1.97 | 0.95 |
| Firewood | (t) | 0.33 | 0.36 | 0.63 | 0.51 | 1.00 | 0.49 |
| Fuel oil | (m ³) | 1.00 | 1.09 | 1.94 | 1.56 | 3.06 | 1.48 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

c) Conversion Factors for Mass


| Multiplied by from | to | kg | t | tl | tc | lb |
|-----------------------|------|-------|----------|----------|----------|---------|
| | | (kg) | (t) | (tl) | (tc) | (lb) |
| Kilogram | (kg) | 1 | 0.001 | 0.000984 | 0.001102 | 2.2046 |
| Metric Ton | (t) | 1,000 | 1 | 0.984 | 1.1023 | 2,204.6 |
| Long ton | (tl) | 1,016 | 1.016 | 1 | 1.12 | 2,240 |
| Short ton | (tc) | 907.2 | 0.9072 | 0.893 | 1 | 2,000 |
| Pound | (lb) | 0.454 | 0.000454 | 0.000446 | 0.0005 | 1 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

d) Conversion Factors for Volume

| Multiplied by from | to | m ³ | l | gal (US) | gal (UK) | bbl | ft ³ |
|-----------------------|----------------------|-------------------|-------|----------|----------|---------|--------------------|
| | | (m ³) | (l) | (US) | (UK) | (bbl) | (ft ³) |
| Cubic meters | (m ³) | 1 | 1,000 | 264.2 | 220 | 6.289 | 35.3147 |
| Liters | (l) | 0.001 | 1 | 0.2642 | 0.22 | 0.0063 | 0.0353 |
| Gallons | (US) | 0.0038 | 3.785 | 1 | 0.8327 | 0.02381 | 0.1337 |
| Gallons | (UK) | 0.0045 | 4.546 | 1.201 | 1 | 0.02859 | 0.1605 |
| Barrels | (bbl) | 0.159 | 159 | 42 | 34.97 | 1 | 5.615 |
| Cub feet | (feet ³) | 0.0283 | 28.3 | 7.48 | 6.229 | 0.1781 | 1 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 34 of 44 |

e) Conversion Factors for Energy

| Multiplied by from | to | J | BTU | cal | kWh |
|--------------------------|-------|-------------------------|--------------------------|------------------------|---------------------------|
| Joule | (J) | 1 | 947.8 x 10 ⁻⁶ | 0.23884 | 277.7 x 10 ⁻⁹ |
| British Thermal Unit | (BTU) | 1.055 x 10 ³ | 1 | 252 | 293.07 x 10 ⁻⁶ |
| Calorie | (cal) | 4.1868 | 3.968 x 10 ⁻³ | 1 | 1.163 x 10 ⁻⁶ |
| Kilowatt-hour | (kWh) | 3.6 x 10 ⁶ | 3412 | 860 x 10 ³ | 1 |
| Ton of oil equivalent | (toe) | 41.87 x 10 ⁹ | 39.68 x 10 ⁶ | 10 x 10 ⁹ | 11.63 x 10 ³ |
| Barrel of oil equivalent | (boe) | 5.95 x 10 ⁹ | 5.63 x 10 ⁶ | 1.42 x 10 ⁹ | 1.65 x 10 ³ |

Source: 2013 National Energy Balance - Ministry of Mines and Energy


f) Mean Coefficients of Equivalence for Gaseous Fuels

| Multiplied by from 1,000 m ³ | to | giga-calorie | toe (10,000 kcal/kg) | boe | tec (7,000 kcal/kg) | giga-joule | million s BTU | megawatt-hour (860 kcal/kWh) |
|--|----|--------------|-------------------------|------|------------------------|------------|---------------|---------------------------------|
| Piped gas Rio de Janeiro | | 3.8 | 0.38 | 2.68 | 0.543 | 15.91 | 15.08 | 4.42 |
| Piped gas São Paulo | | 4.5 | 0.45 | 3.17 | 0.643 | 18.84 | 17.86 | 5.23 |
| Coke oven gas | | 4.3 | 0.43 | 3.03 | 0.614 | 18.00 | 17.06 | 5.00 |
| Dry natural gas | | 8.8 | 0.88 | 6.20 | 1.257 | 36.84 | 34.92 | 10.23 |
| Humid natural gas | | 9.93 | 0.993 | 6.99 | 1.419 | 41.58 | 39.40 | 11.55 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

g) Mean Coefficients of Equivalence for Liquid Fuels

| Multiplied by from m ³ | to | giga-calorie | toe (10,000 kcal/kg) | Boe | tec (7,000 kcal/kg) | giga-joule | millions BTU | megawatt-hour (860 kcal/kWh) |
|--------------------------------------|----|--------------|-------------------------|------|------------------------|------------|--------------|---------------------------------|
| Anhydrous ethyl alcohol | | 5.34 | 0.534 | 3.76 | 0.763 | 22.35 | 21.19 | 6.21 |
| Hydrated ethyl alcohol | | 5.01 | 0.510 | 3.59 | 0.728 | 21.34 | 20.22 | 5.93 |
| Asphalts | | 10.18 | 1.018 | 7.17 | 1.455 | 42.63 | 40.40 | 11.84 |
| Petroleum coke | | 8.73 | 0.873 | 6.15 | 1.247 | 36.53 | 34.62 | 10.15 |
| Refinery gas | | 6.55 | 0.655 | 4.61 | 0.936 | 27.43 | 26.00 | 7.62 |
| Automotive gasoline | | 7.70 | 0.770 | 5.42 | 1.099 | 32.22 | 30.54 | 8.95 |
| Aviation gasoline | | 7.63 | 0.763 | 5.37 | 1.090 | 31.95 | 30.28 | 8.88 |
| LPG | | 6.11 | 0.611 | 4.30 | 0.872 | 25.56 | 24.22 | 7.10 |
| Agents, Lubrication | | 8.91 | 0.891 | 6.27 | 1.272 | 37.29 | 35.34 | 10.36 |
| Naphtha | | 7.65 | 0.765 | 5.39 | 1.093 | 32.05 | 30.37 | 8.90 |


| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 35 of 44 |

| | | | | | | | |
|--|------|-------|------|-------|-------|-------|-------|
| Fuel oil | 9.59 | 0.959 | 6.75 | 1.370 | 40.15 | 38.05 | 11.15 |
| Diesel Oil | 8.48 | 0.848 | 5.97 | 1.212 | 35.52 | 33.66 | 9.87 |
| Other petroleum based energy sources | 8.90 | 0.890 | 6.27 | 1.271 | 37.25 | 35.30 | 10.35 |
| Other non-petroleum based energy sources | 8.90 | 0.890 | 6.27 | 1.271 | 37.25 | 35.30 | 10.35 |
| Petroleum | 8.90 | 0.890 | 6.27 | 1.271 | 37.25 | 35.30 | 10.35 |
| Aviation kerosene | 8.22 | 0.822 | 5.79 | 1.174 | 34.40 | 32.60 | 9.56 |
| Illuminating kerosene | 8.22 | 0.822 | 5.79 | 1.174 | 34.40 | 32.60 | 9.56 |
| Solvents | 7.81 | 0.781 | 5.50 | 1.115 | 32.69 | 30.98 | 9.08 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

h) Mean Coefficients of Equivalence for Solid Fuels


| Multiplied by from ton | to | giga-calorie | toe (10,000 kcal/kg) | boe | tec (7,000 kcal/kg) | giga-joule | millions BTU | megawatt-hour (860 kcal/kWh) |
|-----------------------------|----|--------------|----------------------------|------|---------------------------|------------|-----------------|------------------------------------|
| Tar | | 8.55 | 0.855 | 6.02 | 1.221 | 35.80 | 33.93 | 9.94 |
| Sugarcane bagasse | | 2.13 | 0.213 | 1.50 | 0.304 | 8.92 | 8.45 | 2.48 |
| Guarapa | | 0.62 | 0.062 | 0.44 | 0.089 | 2.61 | 2.47 | 0.72 |
| Imported metallurgical coal | | 7.40 | 0.740 | 5.21 | 1.057 | 30.98 | 29.36 | 8.61 |
| National metallurgical coal | | 6.42 | 0.642 | 4.52 | 0.917 | 26.88 | 25.47 | 7.47 |
| Steam coal 3,100 kcal/kg | | 2.95 | 0.295 | 2.08 | 0.421 | 12.35 | 11.70 | 3.43 |
| Steam coal 3,300 kcal/kg | | 3.10 | 0.310 | 2.18 | 0.443 | 12.98 | 12.30 | 3.61 |
| Steam coal 3,700 kcal/kg | | 3.50 | 0.350 | 2.46 | 0.500 | 14.65 | 13.89 | 4.07 |
| Steam coal 4,200 kcal/kg | | 4.00 | 0.400 | 2.82 | 0.571 | 16.75 | 15.87 | 4.65 |
| Steam coal 4,500 kcal/kg | | 4.25 | 0.425 | 2.99 | 0.607 | 17.79 | 16.86 | 4.94 |
| Steam coal 4,700 kcal/kg | | 4.45 | 0.445 | 3.13 | 0.636 | 18.63 | 17.66 | 5.18 |
| Steam coal 5,900 kcal/kg | | 5.60 | 0.560 | 3.94 | 0.800 | 23.45 | 22.22 | 6.51 |
| Steam coal 6,000 kcal/kg | | 5.70 | 0.570 | 4.01 | 0.814 | 23.86 | 22.62 | 6.63 |
| Unspecified steam coal | | 2.85 | 0.285 | 2.01 | 0.407 | 11.93 | 11.31 | 3.31 |
| Steam coal 5,200 kcal/kg | | 4.90 | 0.490 | 3.45 | 0.700 | 20.52 | 19.44 | 5.70 |
| Charcoal | | 6.46 | 0.646 | 4.55 | 0.923 | 27.05 | 25.63 | 7.51 |
| Mineral coal coke | | 6.90 | 0.690 | 4.86 | 0.986 | 28.89 | 27.38 | 8.02 |
| Firewood | | 3.10 | 0.310 | 2.18 | 0.443 | 12.98 | 12.30 | 3.61 |
| Lye | | 2.86 | 0.286 | 2.01 | 0.409 | 11.97 | 11.35 | 3.33 |
| Molasses | | 1.85 | 0.185 | 1.30 | 0.264 | 7.75 | 7.34 | 2.15 |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 36 of 44 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

i) Densities and Calorific Values – 2012

| Energetic | Density kg/m ³ ⁽¹⁾ | Higher calorific value kcal/kg | Inferior calorific value kcal/kg |
|---------------------------------------|---|-----------------------------------|-------------------------------------|
| Tar | 1,000 | 9,000 | 8,550 |
| Anhydrous ethyl alcohol | 791 | 7,090 | 6,750 |
| Hydrated ethyl alcohol | 809 | 6,650 | 6,300 |
| Asphalts | 1,025 | 10,500 | 9,790 |
| Sugarcane bagasse ¹ | 130 | 2,257 | 2,130 |
| Biodiesel (B100) | 880 | 9,345 | 9,000 |
| Sugarcane juice | - | 623 | 620 |
| Imported metallurgical coal | - | 7,700 | 7,400 |
| National metallurgical coal | - | 6,800 | 6,420 |
| Steam coal 3,100 kcal/kg | - | 3,100 | 2,950 |
| Steam coal 3,300 kcal/kg | - | 3,300 | 3,100 |
| Steam coal 3,700 kcal/kg | - | 3,700 | 3,500 |
| Steam coal 4,200 kcal/kg | - | 4,200 | 4,000 |
| Steam coal 4,500 kcal/kg | - | 4,500 | 4,250 |
| Steam coal 4,700 kcal/kg | - | 4,700 | 4,450 |
| Steam coal 5,200 kcal/kg | - | 5,200 | 4,900 |
| Steam coal 5,900 kcal/kg | - | 5,900 | 5,600 |
| Steam coal 6,000 kcal/kg | - | 6,000 | 5,700 |
| Unspecified steam coal | - | 3,000 | 2,850 |
| Charcoal | 250 | 6,800 | 6,460 |
| Mineral coal coke | 600 | 7,300 | 6,900 |
| Petroleum coke | 1,040 | 8,500 | 8,390 |
| Electricity ² | - | 860 | 860 |
| Hydraulic Power ² | 1,000 | 860 | 860 |
| Piped gas Rio de Janeiro ³ | - | 3,900 | 3,800 |
| Piped gas São Paulo ³ | - | 4,700 | 4,500 |
| Coke oven gas ³ | - | 4,500 | 4,300 |
| Refinery gas | 0.780 | 8,800 | 8,400 |
| Liquefied Petroleum Gas | 552 | 11,750 | 11,100 |
| Dry natural gas ^{3,4} | 0.740 | 9,256 | 8,800 |
| Humid natural gas ^{3,4} | 0.740 | 10,454 | 9,930 |
| Automotive gasoline | 742 | 11,220 | 10,400 |
| Aviation gasoline | 726 | 11,290 | 10,600 |
| Gathered firewood | 300 | 3,300 | 3,100 |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 37 of 44 |

| Energetic | Density kg/m ³ ⁽¹⁾ | Higher calorific value kcal/kg | Inferior calorific value kcal/kg |
|--|---|-----------------------------------|-------------------------------------|
| Commercial firewood | 390 | 3,300 | 3,100 |
| Lye | 1090 | 3,030 | 2,860 |
| Agents, Lubrication | 875 | 10,770 | 10,120 |
| Molasses | 1,420 | 1,930 | 1,850 |
| Naphtha | 702 | 11,320 | 10,630 |
| Fuel oil | 1,000 | 10,085 | 9,590 |
| Diesel Oil | 840 | 10,750 | 10,100 |
| Other petroleum based energy sources | 864 | 10,800 | 10,200 |
| Other non-petroleum based energy sources | 864 | 10,800 | 10,200 |
| Petroleum | 884 | 10,800 | 10,190 |
| Aviation kerosene | 799 | 11,090 | 10,400 |
| Illuminating kerosene | 799 | 11,090 | 10,400 |
| Solvents | 741 | 11,240 | 10,550 |

Source: 2013 National Energy Balance - Ministry of Mines and Energy

1 Bagasse with 50% humidity


2 kcal/kWh

3 kcal/m³


4 At a temperature of 20 °C, for derivatives of petroleum and natural gas

j) Conversion Factors for mean toe

| Energy Source | Unit | toe |
|-----------------------------|----------------|-------|
| Tar | m ³ | 0.855 |
| Anhydrous ethyl alcohol | m ³ | 0.534 |
| Hydrated ethyl alcohol | m ³ | 0.510 |
| Asphalts | m ³ | 1.018 |
| Sugarcane bagasse | T | 0.213 |
| Biodiesel (B100) | m ³ | - |
| Guarapa | T | 0.062 |
| Imported metallurgical coal | T | 0.740 |
| National metallurgical coal | T | 0.642 |
| Steam coal 3,100 kcal/kg | T | 0.295 |
| Steam coal 3,300 kcal/kg | T | 0.310 |
| Steam coal 3,700 kcal/kg | T | 0.350 |
| Steam coal 4,200 kcal/kg | T | 0.400 |
| Steam coal 4,500 kcal/kg | T | 0.425 |

| | | |
|---|--|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 38 of 44 |


| Energy Source | Unit | toe |
|--|--------------------------------|-------|
| Steam coal 4,700 kcal/kg | T | 0.445 |
| Steam coal 5,200 kcal/kg | T | 0.490 |
| Steam coal 5,900 kcal/kg | T | 0.560 |
| Steam coal 6,000 kcal/kg | T | 0.570 |
| Unspecified steam coal | T | 0.285 |
| Charcoal | T | 0.646 |
| Mineral coal coke | T | 0.690 |
| Petroleum coke | m ³ | 0.873 |
| Electricity | MWh | 0.086 |
| Piped gas Rio de Janeiro | 10 ³ m ³ | 0.880 |
| Piped gas São Paulo | 10 ³ m ³ | 0.450 |
| Coke oven gas | 10 ³ m ³ | 0.430 |
| Refinery gas | 10 ³ m ³ | 0.655 |
| Liquefied petroleum gas | m ³ | 0.611 |
| Dry natural gas | 10 ³ m ³ | 0.880 |
| Humid natural gas | 10 ³ m ³ | 0.993 |
| Automotive gasoline | m ³ | 0.770 |
| Aviation gasoline | m ³ | 0.763 |
| Hydraulic | MWh | 0.086 |
| Commercial firewood | T | 0.310 |
| Lye | T | 0.286 |
| Agents, Lubrication | m ³ | 0.891 |
| Molasses | T | 0.185 |
| Naphtha | m ³ | 0.765 |
| Fuel oil (medium) | m ³ | 0.959 |
| Diesel Oil | m ³ | 0.848 |
| Other non-renewable | Toe | 1.000 |
| Other renewable | Toe | 1.000 |
| Other petroleum based energy sources | m ³ | 0.890 |
| Other non-petroleum based energy sources | m ³ | 0.890 |
| Petroleum | m ³ | 0.891 |
| Aviation kerosene | m ³ | 0.822 |
| Illuminating kerosene | m ³ | 0.822 |
| Solvents | m ³ | 0.781 |

| | | |
|---|---|----------------------|
|  | TG01-0.0-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: 0.0 (pilot) |
| | LIFE TECHNICAL GUIDE - 01 Measuring the Biodiversity Pressure Index and definition of Biodiversity Minimum Performance | Page 39 of 44 |

| Energy Source | Unit | toe |
|---------------------------------------|------|--------|
| Uranium contained in UO ₂ | Kg | 73.908 |
| Uranium U ₃ O ₈ | Kg | 10.139 |

Source: 2015 National Energy Balance - Ministry of Mines and Energy

DRAFT

| | | |
|---|---|----------------------|
|  | LIFE-BR-TG01-3.2-English | Revision: 6/1/2021 |
| | Applicability: Europe | Version: Pilot (0.0) |
| | LIFE TECHNICAL GUIDE - 01 Calculation of the Biodiversity Pressure Index and definition of minimum performance in Conservation Actions | Page 40 of 44 |

6. Scopes of GHG Protocol Program

| Sector | Emission Source Scope 1 | Emission Source Scope 2 | Emission Source Scope 3 |
|-----------------------------|--|--|---|
| Energy | | | |
| Generation of Energy | <ul style="list-style-type: none"> Stationary combustion (boilers and turbines used in the production of energy, heat, or steam; fuel pumps; fuel cells; burning of discarded gases or flaring) Mobile combustion (trucks, vessels, and trains for transporting fuels) Fugitive emissions (CH₄ leak from transmission and from storage installations; HFC emissions from storage installations; SF₆ emissions from transmission and distribution equipment) | <ul style="list-style-type: none"> Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> Stationary combustion (mining and extraction of fuels, energy for refining and processing of fuels) Process emissions (productions of fuels, SF₆ emissions) Mobile combustion (transport of fuels / waste, business trips, employee commuting to-from work) Fugitive emissions (CH₄ and CO₂ from landfills, pipelines, SF₆ emissions) |
| Oil & Gas | <ul style="list-style-type: none"> Stationary combustion (process heaters, motors, turbines, burning of discarded gases or flaring, incinerators, oxidants, production of electricity, heat, and steam) Process emissions (process vents, equipment vents, routine and maintenance activities, non-routine activities) Mobile combustion (transport of raw materials, products, waste; vehicles belonging to the company) | <ul style="list-style-type: none"> Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> Stationary combustion (use of products as fuel or combustion for the production of acquired materials) Mobile combustion (transport of raw materials, products, and waste; employees' business trips; employee commuting to-from work; use of products as fuel) Process emissions (use of product as raw material or emissions resulting from the production of acquired materials) Fugitive emissions (CH₄ and CO₂ from landfills or from the production of acquired materials) |

| Sector | Emission Source Scope 1 | Emission Source Scope 2 | Emission Source Scope 3 |
|-------------------------------------|--|--|---|
| | <ul style="list-style-type: none"> Fugitive emissions (leaks from pressurized equipment, sewage treatment, dams) | | |
| Coal Mining | <ul style="list-style-type: none"> Stationary combustion (flaring and use of methane, use of explosives, fires in mines) Mobile combustion (mining equipment, transport of coal) Fugitive emissions (CH₄ emissions from coal mines and coal deposits) | <ul style="list-style-type: none"> Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> Stationary combustion (use of product as fuel) Mobile combustion (transport of coal or waste, employees' business trips, employee commuting to-from work) Process emissions (gasification) |
| Metals | | | |
| Aluminum | <ul style="list-style-type: none"> Stationary combustion (processing of bauxite into aluminum; coke baking; use of lime; sodium carbonate and fuel; CHP) Process emissions (anodic oxidation, electrolysis, PFC) Mobile combustion (transport pre-and post-casting smelting, ore trucks) Fugitive emissions (CH₄, HFC and PFC from fuel pipes, SF₆ as blanket gas) | <ul style="list-style-type: none"> Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> Stationary combustion (processing of raw materials and production of coke by third parties, manufacture of machinery for the production line) Mobile combustion (transport services, business trips, employee's trips) Process emissions (during the production of acquired materials) Fugitive emissions (CH₄ and CO₂ from mining and landfills, emissions from outsourced processes) |
| Iron and steel | <ul style="list-style-type: none"> Stationary combustion (flows of coke, coal, and carbonate; boilers; burners) Process emissions (oxidation of pig-iron, consumption of reducing agent, carbon content of pig-iron and ferroalloys) Mobile combustion (on-site transport) Fugitive emissions (CH₄, N₂O) | <ul style="list-style-type: none"> Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> Stationary combustion (mining equipment, production of acquired materials) Process emissions (production of ferroalloys) Mobile combustion (transport of raw materials, products, waste and intermediary products) Fugitive emissions (CH₄ and CO₂ from sanitary landfills) |
| Chemicals | | | |
| Nitric acid, ammonia, adipic | <ul style="list-style-type: none"> Stationary combustion (boilers, burners, reducing furnaces, flame reactors, steam reformers) | <ul style="list-style-type: none"> Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> Stationary combustion (production of acquired materials, waste combustion) Process emissions (production of acquired materials) |

| Sector | Emission Source Scope 1 | Emission Source Scope 2 | Emission Source Scope 3 |
|--|--|--|--|
| acid, urea, petrochemicals | <ul style="list-style-type: none"> • Process emissions (oxidation or reduction of substrates, removal of impurities, N₂O by-products, catalytic cracking, and several other individual emissions from each process) • Mobile combustion (transport of raw materials, products and waste) • Fugitive emissions (use of HFC, leakage from storage tanks) | | <ul style="list-style-type: none"> • Mobile combustion (transport of raw materials, products and waste; business trips, employee commuting to-from work) • Fugitive emissions (CH₄ and CO₂ from sanitary landfills and ducts) |
| Minerals | | | |
| Cement and lime | <ul style="list-style-type: none"> • Process emissions (calcination of limestone) • Stationary combustion (clinker over, drying of raw materials, energy production) • Mobile combustion (quarry operations, on-site transport) | <ul style="list-style-type: none"> • Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> • Stationary combustion (production of acquired materials, waste combustion) • Process emissions (production of acquired clinker and lime) • Mobile combustion (transport of raw materials, products and waste; business trips, employee commuting to-from work) |
| Waste | | | |
| Landfills, waste combustion, water services | <ul style="list-style-type: none"> • Stationary combustion (incinerators, boilers, burners) • Process emissions (sewage treatment, nitrogen loading) • Fugitive emissions (emissions of CH₄ and CO₂ from the decomposition of waste and animal product) • Mobile combustion (transport of waste or products) | <ul style="list-style-type: none"> • Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> • Stationary combustion (recycled waste used as fuel) • Process emissions (recycled waste used as raw materials) • Mobile combustion (transport of waste or products, business trips, employee commuting to-from work) |
| Pulp & Paper | | | |
| Pulp & Paper | <ul style="list-style-type: none"> • Stationary combustion (production of steam and energy, emissions derived from fossil fuels from the calcination of calcium carbonate in lime ovens, drying | <ul style="list-style-type: none"> • Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> • Stationary combustion (production of acquired materials, waste combustion) • Process emissions (production of acquired materials) |

| Sector | Emission Source Scope 1 | Emission Source Scope 2 | Emission Source Scope 3 |
|---|--|---|--|
| | of products using infrared dryers powered by fossil fuels <ul style="list-style-type: none"> • Mobile combustion (transport of raw materials, products, and waste; operation of harvesting equipment) • Fugitive emissions (CH₄ and CO₂ from waste) | | <ul style="list-style-type: none"> • Mobile combustion (transport of raw materials, products, and waste; business trips, employee commuting to-from work) • Fugitive emissions (landfill emissions of CH₄ and CO₂) |
| Production of HFC, PFC, SF₆ and HCFC-22 | | | |
| Production of HCFC-22 | <ul style="list-style-type: none"> • Stationary combustion (energy consumption, heat or steam) • Process emissions (ventilation of HFC) • Mobile combustion (transport of raw materials, products and waste) • Fugitive emissions (use of HFC) | <ul style="list-style-type: none"> • Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> • Stationary combustion (production of acquired materials) • Process emissions (production of acquired materials) • Mobile combustion (transport of raw materials, products and waste; business trips, employee commuting to-from work) • Fugitive emissions (leakages in the use of the product, CH₄ and CO₂ from landfills) |
| Production of Semiconductors | | | |
| Production of Semiconductors | <ul style="list-style-type: none"> • Process emissions ((C₂F₆, CH₄, CHF₃, SF₆, NF₃, C₃F₈, C₄F₈, N₂O used in the fabrication of wafers, CH₄ created from the processing of C₂F₆ and C₃F₈) • Stationary combustion (oxidation of volatile organic waste; production of energy, heat, or steam) • Fugitive emissions (leakages in the storage of process gases, leakages of remnants from storage tanks) • Mobile combustion (transport of raw materials, products and waste) | <ul style="list-style-type: none"> • Stationary combustion (consumption of energy, heat or steam acquired) | <ul style="list-style-type: none"> • Stationary combustion (production of imported materials, combustion of waste, losses in T&D of energy acquired higher in the value chain) • Process emissions (production of acquired materials, outsourced elimination of gases from processes and remnants from storage tanks) • Mobile combustion (transport of raw materials, products and waste; business trips, employee commuting to-from work) • Fugitive emissions (emissions of CH₄ and CO₂ from landfills, leakages of remnants in storage tanks of process gases lower in the value chain). |
| Other Sectors | | | |

| Sector | Emission Source Scope 1 | Emission Source Scope 2 | Emission Source Scope 3 |
|--|---|--|---|
| Sector of services / organizations with activities performed in offices | <ul style="list-style-type: none"> • Stationary combustion (production of energy, heat or steam) • Mobile combustion (transport of raw materials or waste) • Fugitive emissions (mainly emissions of HFC during the use of refrigeration and air-conditioning equipment) | <ul style="list-style-type: none"> • Stationary combustion (energy consumption, heat or steam acquired) | <ul style="list-style-type: none"> • Stationary combustion (production of acquired materials) • Process emissions (production of acquired materials) • Mobile combustion (transport of raw materials, products and waste; business trips, employee commuting to-from work) |

Source: Specifications of the GHG Protocol Program – 2nd Edition.

DRAFT